

# **FL Peninsular Hendry County - Seminole Big Cypress Reservation Tie Analysis**

and

# **FL Peninsular Hendry County - Palm Beach County Edge Tie Analysis**

SUBMITTED BY:

**Dewberry**

1000 North Ashley Drive Suite 801  
Tampa, FL 33602

*Point of Contact:*

Elise MacPherson, Project Manager  
813.421.8647

SUBMITTED TO:

United States Geological Survey

Table of Contents

Introduction .....3

Edge-Tie Analysis .....3

    Difference Raster .....3

    Elevation Difference Thresholds.....5

Edge-Tie Results .....6

    Seminole Big Cypress Reservation.....6

Vegetation & Water Levels .....6

    Palm Beach County AOI .....7

    Water Levels.....8

Summary .....9

## Introduction

Dewberry was tasked to evaluate how well the newly produced FL Peninsular Hendry County (acquired in late 2018 to early 2019) ties spatially to preexisting lidar data produced for the Seminole Big Cypress Reservation project and the Palm Beach County lidar project. The lidar data acquired for the Seminole Big Cypress Reservation project was originally acquired for USGS in 2014. The lidar data acquired for the Palm Beach County project was originally acquired for USGS in 2016. Dewberry has compared the new Hendry County lidar data to the existing Seminole Big Cypress data & the existing Palm Beach County data where the two datasets overlap.

## Edge-Tie Analysis

There are 30 Hendry tiles which overlap with the Seminole Big Cypress Reservation lidar data, there are 34 Hendry tiles which overlap with the Palm Beach County lidar data, shown in Figure 1 below. A difference raster was produced to analyze elevation differences between the two datasets in areas of overlap. Profiles and visual reviews were used to compare the two datasets where the datasets are adjacent, but do not overlap. This review of the adjacent, non-overlapping areas was to ensure no obvious feature discontinuities exist between the datasets. Dewberry has determined that no gross feature discontinuities were identified.




Figure 1- Thirty tiles overlap between the Hendry AOI and previously collected Seminole Big Cypress Reservation AOI, there are 34 tiles that overlap the Hendry AOI and the Palm Beach County AOI.

## DIFFERENCE RASTER

The USGS Seminole Big Cypress Reservation DEMs match the 2.5 ft DEM size specified for Hendry county, the data was consistent. Using the 2.5 ft bare-earth DEMs for each dataset, Dewberry created a

difference raster by subtracting Seminole Big Cypress Reservation data from Hendry county data (Hendry-Seminole Big Cypress Reservation). This difference raster is shown in Figure 2 below.




Figure 2-Difference raster and symbology key created for the Hendry- Seminole Big Cypress Reservation overlap.

The USGS Palm Beach County DEMs match the 2.5 ft DEM size specified for Hendry county, the data was consistent. Using the 2.5 ft bare-earth DEMs for each dataset, Dewberry created a difference raster by subtracting Palm Beach County data from Hendry county data (Hendry-Palm Beach County). This difference raster is shown in Figure 3 below.




Figure 3-Difference raster and symbology key created for the Hendry- Palm Beach County overlap.

All hydrographic features breaklined in Hendry, Seminole Big Cypress Reservation, and Palm Beach County including streams, rivers, ponds, and lakes were excluded from analysis as water levels varied between the two lidar acquisitions. Hydrographic features, overlaid on the difference raster, are shown in blue in Figure 4 below.




Figure 4-Breaklined hydrographic, shown in blue fill, were removed from statistical analysis generated from the difference raster as water levels may vary between the different lidar acquisition years. This example is from the Palm Beach County data.

### **ELEVATION DIFFERENCE THRESHOLDS**

Per client discussions and requests, all overlapping data should be analyzed to show which areas have elevations within 10 cm (RMSEz requirements for USGS QL1 and QL2 data) of each other. As such, pixels in the difference raster representing 0 to +10 cm and 0 to -10 cm of elevation change between the two datasets are binned. From these initial bins, Dewberry then used thresholds of 20 cm up until +/-90 cm as this allows detailed analysis of changes occurring which are less than +/-1 m in difference. The 20 cm bins, starting from the required +/-10 cm bin also allows for analysis of change at +/-30 cm, which is the required VVA for USGS QL1 and QL2 data. Larger elevation differences tend to result from similar or consistent sources, so after the +/-90 cm bins, data are binned to +/-2.5 m, +/-5 m, and everything greater than +5 m or less than - 5 m. If the units of data are in feet, the metric values listed above are converted to feet for analysis.

Dewberry symbolized the difference rasters for this analysis using the binned values and color schema shown below. Values are in feet.




Figure 5- Pixels within the 0 to +/-30 cm (0.98 ft) threshold are colored as green. Dark green is used for pixels in the 0 to +/- 10 cm (0.33 ft) bin and light green is used for the +/-10 cm (0.33 ft) to +/- 30 cm (0.98 ft) bin. The Hendry Seminole Big Cypress Reservation symbology is on the left and on the right is the Hendry Palm Beach difference raster.

## EDGE-TIE RESULTS

### Seminole Big Cypress Reservation

When looking at all overlap areas consisting of all slopes and all land cover types, 38.87% of the overlapping points are within the 0 to +/-10 cm (0.33 ft) threshold with the majority of these points being located in flat, open terrain. Additional analysis shows 77.77% of the overlapping points are within the 0 to +/-30 cm (0.98 ft) threshold with the majority of these points being located in vegetated areas. These variations are allowable elevation differences between the two datasets.

The remaining points that exceed +/-30 cm (0.98 ft) are located in areas with temporal offsets. The temporal offsets may occur in dynamic, non-static environments, such as wetlands, marsh, or floodplains, or may occur in less dynamic, upland areas due to man-made or cultural changes. The temporal offset cause in this overlap area are located in wetland/marsh areas. The differences greater than +/- 30 cm (0.98 ft) are described in the sections below.

### VEGETATION & WATER LEVELS

The majority of areas with larger vertical differences between these two datasets occur within vegetated areas and areas with changing water levels. These types of changes are due to the ~5 year temporal difference between the two lidar acquisitions.

The figures below show examples of these temporal changes.




Figure 6- The image above shows an area in the difference raster and basemap imagery with a temporal difference in the dynamic, non-static environments (wetland marsh) as well as water levels where the areas were not breaklined due to the specification requirements.




Figure 7- The image above shows another area in the difference raster and basemap imagery with a temporal difference in the dynamic, non-static environments (wetland marsh and crop fields) as well as water levels where the areas were not breaklined due to the specification requirements.

#### Palm Beach County AOI

When looking at all overlap areas consisting of all slopes and all land cover types, 39.89% of the overlapping points are within the 0 to +10 cm (0.33 ft) threshold with the majority of these points being located in flat, open terrain. Additional analysis shows 76.16% of the overlapping points are within the 0

to +/-30 cm (0.98 ft) threshold with the majority of these points being located in vegetated areas. These variations are allowable elevation differences between the two datasets.

The remaining points that exceed +/-30 cm (0.98 ft) are located in areas with temporal offsets. The temporal offsets may occur in dynamic, non-static environments, such as wetlands, marsh, or floodplains, or may occur in less dynamic, upland areas due to man-made or cultural changes.. The temporal offset cause in this overlap area are located in wetland/marsh areas. The differences greater than +/- 30 cm (0.98 ft) are described in the sections below.

### **WATER LEVELS**

The majority of areas with larger vertical differences between these two datasets occur within vegetated areas and areas with changing water levels. These types of changes are due to the ~3 year temporal difference between the two lidar acquisitions.

The figure below show examples of these temporal changes.




Figure 8- The image above shows an area in the difference raster and basemap imagery with a temporal difference in the dynamic, non-static environments (crop field).



## Summary

Overall the Hendry and Seminole Big Cypress Reservation lidar data match well with 38.87% of the overlap data matching within 0 to +/-10 cm (0.33 ft) and 77.77% of the overlap data matching within +/-30 cm (0.98 ft) . The areas of largest vertical elevation change occur due to temporal differences and include varying levels of water in hydrographic features and changes in wetlands/marsh areas. Cultural or man-made changes also contribute to larger elevation differences, including new hydrographic control structures, new reservoirs or impoundments, and likely construction/roadway improvements.

The Hendry and Palm Beach County lidar data match well with 39.89% of the overlap data matching within 0 to +/-10 cm (0.33 ft) and 76.16% of the overlap data matching within +/-30 cm (0.98 ft) . The areas of largest vertical elevation change occur due to temporal differences and include varying levels of water in hydrographic features and changes in wetlands/marsh areas. Cultural or man-made changes also contribute to larger elevation differences, including new hydrographic control structures, new reservoirs or impoundments, and likely construction/roadway improvements.